

## ON THE WORST-CASE CONVERGENCE OF MR AND CG FOR SYMMETRIC POSITIVE DEFINITE TRIDIAGONAL TOEPLITZ MATRICES\*

JÖRG LIESEN<sup>†</sup> AND PETR TICHÝ<sup>†</sup>

**Abstract.** We study the convergence of the minimal residual (MR) and the conjugate gradient (CG) method when applied to linear algebraic systems with symmetric positive definite tridiagonal Toeplitz matrices. Such systems arise, for example, from the discretization of one-dimensional reaction-diffusion equations with Dirichlet boundary conditions. Based on our previous results in [J. Liesen and P. Tichý, *BIT*, 44 (2004), pp. 79–98], we concentrate on the next-to-last iteration step, and determine the initial residuals and initial errors for the MR and CG method, respectively, that lead to the slowest possible convergence. By this we mean that the methods have made the least possible progress in the next-to-last iteration step. Using these worst-case initial vectors, we discuss which source term and boundary condition in the underlying reaction-diffusion equation are the worst in the sense that they lead to the worst-case initial vectors for the MR and CG methods. Moreover, we determine (or very tightly estimate) the worst-case convergence quantities in the next-to-last step, and compare these to the convergence quantities obtained from average (or unbiased) initial vectors. The spectral structure of the considered matrices allows us to apply our worst-case results for the next-to-last step to derive worst-case bounds also for other iteration steps. We present a comparison of the worst-case convergence quantities with the classical convergence bound based on the condition number of  $A$ , and finally we discuss the MR and CG convergence for the special case of the one-dimensional Poisson equation with Dirichlet boundary conditions.

**Key words.** Krylov subspace methods, conjugate gradient method, minimal residual method, convergence analysis, tridiagonal Toeplitz matrices, Poisson equation

**AMS subject classifications.** 15A09, 65F10, 65F20

---

\*Received January 15, 2005. Accepted for publication July 11, 2005. Recommended by D. Szyld.

<sup>†</sup>Institute of Mathematics, Technical University of Berlin, Straße des 17. Juni 136, 10623 Berlin, Germany. (liesen@math.tu-berlin.de and tichy@math.tu-berlin.de). This research was supported by the Emmy Noether-Programm of the Deutsche Forschungsgemeinschaft. The work of the second author was also supported by the Grant Agency of Academy of Sciences of the Czech Republic under grant No. KJB1030306.